

Sedimentology and Stratigraphic Architecture of Quaternary Paleochannels Along the Inner Shelf of South Carolina

Joshua H. Long¹ (jhlong@coastal.edu), Till J.J. Hanebuth¹, Clark A. Alexander², Katherine Luciano³

1. Department of Coastal and Marine Systems Science, Coastal Carolina University, Conway, South Carolina; 2. Skidaway Institute of Oceanography, University of Georgia, Savannah, Georgia;

3. South Carolina Department of Natural Resources Geological Survey, Charleston, South Carolina





(Abstract)

Throughout the Quaternary, numerous high frequency, high amplitude glacioeustatic sea level cycles have left behind a complex network of paleochannels and paleovalleys across the inner continental shelf off the coast of South Carolina. These isolated paleotopographic lows were carved into a substrate of pre-Quaternary rocks and Pleistocene deposits by coastal plain and Piedmont-draining rivers and provided rare accommodation in an overall low-accommodation setting. The distribution, age, composition, and stratigraphic architecture of these sedimentary successions offer insight into both the allogenic and autogenic forces that influenced depositional processes at the time of formation.

As part of a 2015 regional sand resource assessment, a dense grid of high resolution subbottom (Chirp) data was acquired off the coast of Folly Beach, South Carolina by the Bureau of Ocean Energy Management (BOEM). These data provide the spatial resolution required to reconstruct paleochannel connectivity and define the internal stratigraphic architecture of paleochannel and paleovalley successions. Paleochannel fills display concentric, asymmetrical, or horizontal geometries reflecting varying degrees of erosion, aggradation, and accretion. Paleovalleys exhibit vertical relief of up to 10 meters and apparent widths of up to 3 kilometers. Within paleovalleys, individual channels are arranged in both multistory and multilateral configurations that are the result of changes in localized autogenic processes as well as regional allogenic controls that influenced accommodation.

In addition to geophysical data, 52 meters of sediment core recovered from 9 locations provide the control necessary to reconstruct depositional processes associated with this architecture. Detailed physical descriptions including sediment composition, lithology, and physical structures; geochemical analysis; ¹⁴C dating; and AAR age estimates of cored intervals reveal channel fill that was deposited during the mid-Holocene and was dominated by tidally-influenced processes in low to mixed energy, estuarine and backbarrier settings.

(Seismic Facies)

Siesmo-stratigraphic Facies			External Bounding	Geosiesmic Color	Cross-sectional Geo	metry	
Primary Secondary		Surfaces Order	Internal		External	Interpretation	
	Fill Type	Concentric (c)	4, 5, 6	СНс	High to moderate amplitude, aggradational, concentric reflectors	Broadly Lenticular	Mud-rich, aggradational fill. Commonly estuarine.
Paleochannel Facies		Horizontal (h)		CHh	High to moderate amplitude, horizontal to sub- horizontal reflectors		Passively filled, likley fine-grained channel fill.
		Asymmetric (a)		Cha	High to moderate amplitude, uniformally-dipping reflectors		Point bar deposits of meandering fluvial or estuarine channels.
Раївос		Chaotic (k)		CHk	Disorganized, moderate amplitude, discontinous reflectors		High energy channel deposits
		Transparent (t)		CHt	Low amplitude homogeneous fill		Homogeneous channel fill
Horizontally Stratified Facies		5 or 6	HS	High to moderate amplitude, horizontal to sub- horizontal reflectors	Highly irregular with erosional upper surface	Fine-grained, non-channelized back barrier or shelf deposits	
Indistinct Facies		6	ID	transparent to chaotic	any	Transgressive sand sheets and other deposits	
Pre-Quaternary Basement			6	PQ	Variable. Transparent to high amplitude horizontal to gently dipping reflectors	Highly irregular, erosional upper surface. Base is unresolvable	Cretaceous through Pliocene consolidated to lithified sediment of variable lithology.

Table 1. Stratigraphic architecture of paleochannels and shelf deposits based on seismic and core data. Interpretations based on observation as well as concepts from Mitchum et al. (1977) and Gibling (2006).

a	Reflector	Configuration			b Succession - dominated	Erosion - dominated	c		
Patterns	Interpretation	Frequency	Amplitude	Continuity			MFS J	Open marine	Shelf
Parallel draped	Deposition under low energy sedimentary regime	A measure of the spacing between	A measure of velocity density contrast	Continuous			SB		
Sub-parallel	Aggradation under relatively uniform energy conditions	reflectors High	High	Continuity in bedding strata or depositional processes	Multilateral	Multistory		Estuarine	Salt
Sigmoid to oblique	Progradation into deeper water	Thin bedding or high variability	Large changes in density of sediment indicating mixed	Discontinuous Discontinuity in bedding strata				Tidal bars	* *
Parallel oblique	Lateral progradation	Medium	Medium	or deposition under a highly variable energy regime Terminations	SINGLE-STORY CHANNEL B	DIES AND INDIVIDUAL STORIES	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	Estuary	* *
Hummocky to oblique	Rapid channel infill under variable energy sedimentary regimes	Intermediate between high and low	Intermediate between high and low	Onlap Aggradation of younger sediments onto a surface	FÖRM	FILL	5	Tidal-Fluvial Channel	
Hummocky to chaotic	Deposition in a variable high energy setting	Low	LOW Small changes in	Downlap Progradation of younger				Fluvial	
Undifferentiated	Homogeneous or highly contorted, steeply dipping strata	Thicker bedding or low variability	density of sediment indicating relative	sediments onto a surface Truncation	Symmetric Asymmetric	Concentric Asymmetric			

Figure 1. A) Basic stratal geometries as summarized in Mellet et al. (2013) from concepts defined by Mitchum et al. (1977) which form the basis for the seismo-stratigraphic facies used in this study. B) Paleochannel and paleovalley architecture from Gibling (2006). C) Conceptual model of tide-dominated estuarine fill within an incised valley displaying the prominant tripartite sequence stratigraphic interpretation defined by lower fluvial LST (Lowstand Systems Tract), intermediate estuarine TST (Transgressive Systems Tract), and upper marine HST (Highstand Systems Tract) deposits. From Dalrymple and Choi (2007) based on Dalrymple (1992).





Figure 2. Base map showing location of seismic and core data used in this study, general distribution of paleoincisions, interpreted paleovalley trends, and relevant previous studies.



Preliminary Results

1) Paleochannel fill is predominantly mud rich and heterolithic, containing sedimentological and paleontological indicators of having been deposited in low-energy, tidally-influenced, brackish water settings.

2) The youngest generation of paleochannels in this area were infilled during the mid-Holocene (~4 - 8 ka) following the stabaliziation of sea level.

3) Mud-rich paleochannel fills are a significnat component of the local geologic framework and are commonly exposed on the modern seafloor, potentially influencing local morphodynamics and serve as a source of fine-grained sediment.

4) The most common types of paleochannel fill in this area are concentric or horizontal, which, along with observations from sediment cores, suggests systems—dominated by suspension sedimentation.

5) Most larger paleochannels (> 1km) occur as part of paleovalley systems rather than in isolation which may suggest a stratigraphic control on drainage network development.

6) Individual paleovalleys can be correlated throughout the data set based on paleochannel architecture and paleochannel associations within each paleovalley.

7) Depth of incision is typically limited to the depth of Pleistocene sediment defined by the top of the pre-Quaternary section (Paleocene or Miocene in this area as defined by Weems et al., 2014 and Harris et al., 2005).

References

Blum, M., Martin, J., Millikin, K., and Garvin, M., 2013. Paleovalley Systems: Insights from Quaternary Analogues and Experiments. Earth-Science Reviews, v. 116, p. 128-169.

Dalrymple, R.W., 1992. Tidal depositional systems. in Walker, R.G., James, N.P. (Eds.), Facies Models: Response to Sea Level Change. Geological Association of Canada, St. John's, pp. 195–218.

Dalrymple, R.W. and Choi, K., 2007. Morphologic and facies trends through the fluvial–marine transition in tide-dominated depositional systems: A schematic framework for environmental and sequence-stratigraphic interpretation. Earth Science Reviews, v. 81, p. 135-174.

Desjardins, P.R., Buatois, L.A., and Mangano, G.M., 2012. Tidal Flats and Subtidal Sand Bodies in Developments in Sedimentology, v. 64. Elsevier publishing, p. 529-561.

Feldman, H., and Demko, T., 2013. Recognition and prediction of petroleum reservoirs in the fluvial/tidal transition in Ashworth, P.J., Best, J.L., and Parsons, D.R., ed. Fluvial Tidal -SedimentologyDevelopments in Sedimentology: Developments in Sedimentology, v. 68, p. 483-528.

Gibling, M.R., 2006. Width and Thickness of Fluvial Channel Bodies and Valley Fills in the Geological Record: A literature Compilation and Classification. Journal of Sedimentary Research, v. 76, p. 731-770.

Harris, S.M., Gayes, P.T., Kindinger, J.L., Flocks, J.G., Krantz, D.E., and Donovan, P., 2005. Quaternary Geomorphology and Modern Coastal Development in Response to an Inherent Geologic Framework: An Example from Charleston, South Carolina. Journal of Coastal Research, v. 21, no. 1, p. 49-64.

Mellet, C.L., Hodgson, D.M., Platera, A.J., Mauz, A., Selby, I., and Lang, A., 2013. Denudation of the continental shelf between Britain and France at the glacial-interglacial timescale. Geomorphology, v. 203, p. 79-96.

Mitchum, R.M., Vail, P. R., and Sangree, J.B.,1977. Seismic Stratigraphy and Global Changes of Sea Level, Part 6: Stratigraphic Interpretation of Seismic Reflection Patterns in Depositional Systems in Payton, C.E., ed., Seismic Stratigraphy - Applications to Hydrocarbon Exploration: AAPG Memoir 26, p. 117-144.

Weems, R. E., Lewis, W. C., Lemon, Earl M., 2014. Surficial Geologic Map of the Charleston Region, Berkeley, Charleston, Colleton, Dorchester, and Georgetown Counties, South Carolina. USGS Open File Report 2013-1030.

Acknowledgement

This work benefitted significantly from financial support provided by the South Carolina Gological Survey and Coastal Carolina University as well as technical input from various contributors involved in the ongoing southeast regional Bureau of Ocean Energy Management (BOEM)Atlantic Sand and Gravel Assessment Project (ASAP).

lithologies and sedimentary structures indicative of tidal influence including alternating mud-sand couplets, thin-thick bedding, mud drapes, and

limited bioturbation (Dalrymple and Choi, 2007; Desjardins et al., 2012; Feldman and Demko, 2013;) as well as shells of intertidal species including

brown mud snails (Illyanasa obseleta), dwarf surf clams (Mulinia lateralis), oysters (Burg, E., Harding, J., and Kelley, P., 2019, personal communications)