

Miocene Tectonic and Stratigraphic Evolution of the Central Baja California Peninsula and Rapid Marine Flooding Into the Gulf of California at ca. 6.3 Ma




Rebecca Dorsey (University of Oregon)





Darin, M.H., Bennett, S.E.K., Hausback, B.P., Gardner, K.K., Niemi, T.M., Busby, C., Graettinger, A., Salgado Muñoz, V.O., Martinez Gutierrez, G., Morebeck, C., Usher, E., Heizler, M.T., Pecha, M.E., Stelten, M., Schmitt, A., Dolby, G.A.

Two Collaborative Projects

Baja Geogenomics (BGG)



Baja Basins (2017 – 2022)




Tectonic Setting

- Gulf of California: opens by relative plate motion across oblique **transform-rift Pac-NAM** plate boundary (long transform faults).
- Baja California Microplate moves NW relative to North America at modern rate (gps): ~45 mm/yr (Plattner et al., 2015, JGR).

Late Cenozoic Tectonic History of Central Baja Peninsula

- 1) 30 – 12 Ma: Farallon plate subduction & arc magmatism
- 2) 12 – 8 Ma: End of subduction, reorganization, oblique extension
- 3) ~ 8 – 0 Ma: Strike-slip motion on plate boundary (in Gulf of Cal.)
 - **Santa Rosalía Basin** - transtensional pull-apart basin
 - Stratigraphic record of rapid **marine flooding at ~ 6.3 Ma** ...
 - During onset of seafloor spreading in Guaymas basin:
 - **Geologically instantaneous** regional marine inundation.






Geology of the Central Baja California Peninsula

- Miocene subduction-related volcanic/volcaniclastic rocks.
- Plio-Pleis. volc & sed deposits
- Belt of strike-slip & normal faults and fault-bounded basins in NE.

Santa Rosalía Basin

- 80-km long, late Mioc. to Pleist. sedimentary rocks.
- Records opening of the Gulf of California transform-rift.
- Initiated ~ 6.3 Ma (this study).
- Inverted/uplifted by La Reforma volcano starting ~ 1.4 Ma.





1

2

3

4



5

6





7

Chronostratigraphy

3 Main Stages




Stages 1 and 2: Tectonic-Magmatic Evolution (Highlights)




M. Heizler

Esperanza Basalt

- Regionally extensive tholeiitic basalt, thin flows, early post-subduction.
- Previously published ages: 7.6 – 12.5 Ma.
- New paleomag and age data (this study): Esperanza has **normal** polarity.
- Basalt emplaced **ca. 9.9 Ma**



(3) Initiation of the Central Gulf of California



 Pleistocene marine terraces, ages ca. 100 ka to ~1.4 Ma (Ortlieb, 1991; Schmitt et al., 2006)

 Pleistocene terraces of uncertain age

QTs Paleocene to Holocene sedimentary deposits (includes San Gregorio, Isidro, Ignacio Fms)

 Pv Pleistocene calc-alkaline volcanic rocks

Pb Plio-Pleistocene magnesian andesites and basalts

Mb Late Miocene (~12-9 Ma) post-subduction tholeiitic lavas, Nb-enriched basalts, adakites, and magnesian andesites

Mv Comondú (early-mid Miocene) subduction-related volcanic rocks AND post-Comondú (late Mioc.) andesitic to basaltic rocks

Pz-Mz Paleozoic and Mesozoic plutonic and metamorphic rocks

Geology of the Central Baja California Peninsula

- Miocene subduction-related volcanic/volcaniclastic rocks.
- Plio-Pleis. volc & sed deposits
- Belt of strike-slip & normal faults and fault-bounded basins in NE.

Santa Rosalía Basin

- 80-km long, late Mioc. to Pleist. sedimentary rocks.
- Latest Miocene Boleo Fm: initial marine flooding ~ **6.3 Ma**.
- Inverted/uplifted by La Reforma volcano starting ~ 1.4 Ma.

Boleo Formation: Sedimentary Facies Associations

Facies Associations

4. Mantos (shale): stratiform ore deposits

Rapid marine transgressions, hydrothermal activity, liquefaction.

3. Gypsum Facies: subaqueous & subaerial


Distal evaporite basin, subaqueous to subaerial, age-equivalent to deltas.

2. Siliciclastic Facies: C-up parasequences

Large input of sediment due to growth of topography in Baja peninsula.

1. Basal Limestone (~ 5 m thick)


Initial marine incursion into Gulf of California, localization of strain along the Pac.-Nam plate boundary.




Boleo Formation: Sedimentary Facies Associations

Marine Basal Limestone (thin, transgressive)

Travertine/Tufa/Stromatolites
Carbonate Shoreline







Calcareenite/Coquina
Nearshore Shelf



Molluscan Packstone
Offshore Shelf




Siliciclastic Facies

	Code	Lithology	Summary Description	Interpret -ation
Coarsening-Up (delta progradation)	S1		Thin Shale Manto	Marine Flooding
	S5		Conglomerate	Alluvial Fan
	S4		Interbedded Congl and Sst	Fan Delta Front
	S3		Sandstone (X-bedded, bioturbated)	Saline Inner Shelf
	S2		Heterolithic Facies	Saline Outer Shelf
	S1		Thin Shale Manto	Marine Flooding
	S5		Conglom.	Alluvial Fan

MFS = marine flooding surface

Gypsum Facies (Salgado Muñoz and Niemi, in review)



	Code	Lithology	Summary Description	Interpret -ation
Shallowing-Up (basin desiccation)	E1		Laminated Gypsum	Deep Evap. Basin
	E5		Gypsrudite (breccia)	Subaerial Exposure
	E4		Mud-Gypsum Laminae	Subaerial Sabkha Flat
	E3		Wavy Laminated Gypsum	Semi-Emergent Shoals
	E2		Laminated Gypsum with Selenite Xls	Shallow Evap. Basin
	E1		Laminated Gypsum	Deep Evap. Basin
	E5		Conglom.	Alluvial Fan

MFS = marine flooding surface



Boleo Formation: Sedimentary Facies Associations

SILICICLASTIC Facies


Mud-silt-sand heterolithic facies *Saline Outer Shelf*



Cross-Bedded and Bioturbated Sst *Saline Inner Shelf*





Channelized Conglomerate & Sst *Alluvial Fan & Fan Delta*



Boleo Formation: Sedimentary Facies Associations






GYPSUM Facies




Thin Shale MANTOS (marine)

Mantos contain marine calcareous nannofossils


Stratiform Cu-Co-Zn ore deposits







PANEL 1

Boleo Formation




PANEL 2

Boleo Formation: Arroyo Infierno





Clast imbrications:
Gravel transport to NE



Geologic-Stratigraphic Cross Section: Arroyo Infierno (northern basin)




Syn-Depositional Growth Strata



Growth Faults in Boleo Fm: Record of syn-depositional faulting and tilting to NE






Monocline Growth: driver of syn-depositional faults


(A) Initiation of monocline above tip of oblique normal fault, marine flooding into subsiding basin, deposition of Boleo marine basal limestone.

B. Monocline growth, faulting, tilting



(B) Growth of monocline above tip of propagating normal fault, NE tilting on antithetic normal faults, continued subsidence, deposition of Boleo Formation siliciclastic (silic.) and gypsum (gyp) facies

C. Main fault breaks to surface




(C) Main fault breaks to the surface, Tirabuzón Formation is deposited on top of older antithetic normal faults.

(adapted from Willsey et al., 2002; Jackson et al., 2006; Wilson et al., 2009).


Transtensional pull-apart basin. Boleo Fm: two-stage response to episodic faulting

Stage 1: Transtensional faulting & rapid marine flooding, transgression.

Mantos record hydrothermal activity.




Stage 2: Basin is isolated from the ocean. Evaporative drawdown & progradation of alluvial fans and fan deltas into the basin.



Basin Model (pull-apart basin)



Tectonic Reconstruction (modif. from Miller & Lizarralde, 2013)

~ 6.3 Ma



Guaymas – Santa Rosalía Pull-Apart Basin during **onset of sea floor spreading**.

Geologically Instantaneous marine flooding into Gulf of California at ca. 6.3 Ma



Marine flooding into pre-existing basins: geologically instantaneous event (within age uncertainties) that inundated a >1,000 km lowland along the plate boundary from the central to northern Gulf of California and Salton Trough to the lower Colorado River valley in as little as ca. 100,000 years.

CONCLUSIONS

Late Cenozoic Tectonic History, Central Baja Peninsula – Three stages:

- 1) 30 – 12 Ma: Farallon plate subduction & arc magmatism (Comondú Gp)
- 2) 12 – 8 Ma: End of subduction, tectonic reorganization, oblique extension
- 3) ~ 8 – 0 Ma: Strike-slip motion in modern Gulf of California
 - **Latest Miocene Boleo Formation (Santa Rosalía Basin):**
 - Transtensional pull-apart basin
 - Mixed evaporite-siliciclastic sedimentation
 - Thin shale mantos: episodic marine flooding, seismicity, hydrothermal activity
 - Boleo Basal limestone records rapid **marine flooding at ~ 6.3 Ma**, during onset of seafloor spreading in Guaymas basin.
 - **Geologically “instantaneous”** marine flooding event inundated >1,000 km of plate boundary lowland in as little as ~ 100,000 years.